

Melatrin cream

Fluocinolone Acetonide USP, Hydroquinone USP & Tretinoin USP

Composition

Melatrin Cream: Fluocinolone Acetonide USP 0.01%, Hydroquinone USP 4% and Tretinoin USP 0.05% in a hydrophilic cream base for topical application.

Pharmacology

Melatrin Cream contains three active ingredients Fluocinolone Acetonide, Hydroquinone and Tretinoin. All of them are used to treat melasma. However, the mechanism of action of the active ingredients in Melatrin Cream in the treatment of melasma is unknown but Fluocinolone Acetonide is a corticosteroid for topical dermatological use and is classified therapeutically as an anti-inflammatory. Hydroquinone is classified therapeutically as a depigmenting agent, which may interrupt one or more steps in the tyrosine-tyrosinase pathway of melanin synthesis. Tretinoin is classified therapeutically as a keratolytic agent.

Indications and Uses

Melatrin Cream is indicated for the short-term treatment of moderate to severe melasma of the face, after starting the treatment avoid presence of sun. Use a sunscreen of at least SPF 30 or more.

Dosage and Administration

Melatrin Cream should be applied once daily at night. It should be applied at least 30 minutes before bedtime. Gently wash the face and neck with a mild cleanser. Rinse and pat the skin dry. Apply a thin film of the cream to the hyperpigmented areas of melasma including about 1/2 inch of normal appearing skin surrounding each lesion. Rub lightly and uniformly into the skin. During the day, use a sunscreen and wear protective clothing. Avoid sunlight exposure. Patients may use moisturizers and/or cosmetics during the day.

Contraindications

Melatrin Cream is contraindicated in individuals with a history of hypersensitivity, allergy or intolerance to this product or any of its components.

Precautions

Melatrin Cream contains Hydroquinone and Tretinoin that may cause mild to moderate irritation. Local irritation, such as skin reddening, peeling, mild burning sensation, dryness and pruritus may be expected at the site of application. Transient skin reddening or mild burning sensation does not preclude treatment. If a reaction suggests hypersensitivity or chemical irritation, the use of the medication should be discontinued. Melatrin Cream also contains the corticosteroid Fluocinolone Acetonide. Systemic absorption of topical corticosteroids can produce reversible Hypothalamic-Pituitary-Adrenal (HPA) axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal of treatment. Manifestations of Cushing's syndrome, hyperglycemia and glucosuria can also be produced by systemic absorption of topical corticosteroid while on treatment. If HPA axis suppression is noted, the use of Melatrin Cream should be discontinued. Recovery of HPA axis function generally occurs upon discontinuation of topical corticosteroids.

Side-effects

A very few patients may get severe allergic reactions from Melatrin Cream. They may have trouble breathing or severe asthma attacks. While patients use Melatrin Cream, skin may develop mild to moderate redness, peeling, burning, dryness or itching.

Drug Interactions

Avoid use of medicated or abrasive soaps, cleansers, soaps, cosmetics with drying effects, products with high concentration of alcohol, astringent & other irritants or keratolytic drugs. Also avoid concomitant use of medications with photosensitizing effects.

Pregnancy & Lactation

Pregnancy: Pregnancy Category C: Melatrin Cream contains the teratogen, tretinoin, which may cause embryo-fetal death, altered fetal growth, congenital malformations and potential neurologic deficits. It is difficult to interpret the animal studies on teratogenicity with Melatrin Cream, because the availability of the dermal applications in these studies cannot be assured and comparison with clinical dosing is not possible. There are no adequate and well-controlled studies in pregnant women. Melatrin Cream should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. **Lactation:** Corticosteroids, when systemically administered, the drug appear in human milk. It is not known whether topical application of Melatrin Cream could result in sufficient systemic absorption to produce detectable quantities of Fluocinolone Acetonide, Hydroquinone or Tretinoin in human milk. Because many drugs are secreted in human milk, caution should be exercised when Melatrin Cream is administered to nursing women.

Pediatric Use : Safety and effectiveness of Melatrin Cream in pediatric patients have not been established.

Storage

Do not store above 30°C. Keep in a dry place & protect from light.

Melatrin Cream 30 g : Each carton contains a tube of 30 g cream, an insert and an usage guideline.

Melatrin Cream 15 g : Each carton contains a tube of 15 g cream, an insert and an usage guideline.

Manufactured by:



Ziska Pharmaceuticals Ltd.
Kaliakoir, Gazipur, Bangladesh