

# Pamix

Paracetamol BP

**Composition:**

**Pamix Tablet:** Each tablet contains Paracetamol BP 500 mg.

**Pamix Suspension:** Each 5 ml suspension contains Paracetamol BP 120 mg.

**Pamix Paediatric drops:** Each ml drop contains Paracetamol BP 80 mg.

**Pharmacology:**

**Pamix** contains paracetamol which is an analgesic and antipyretic agents, specially suitable for patients who cannot tolerate aspirin or other analgesics. Paracetamol inhibits the synthesis of prostaglandin from arachidonic acid by COX-3 pathway.

**Indication:**

All conditions requiring relief from pain, fever due to neuritis, neuralgia, headache, migrant, toothache, rheumatic disorder, cold, influenza, dysmenorrhea, acute & chronic inflammatory disorder.

**Dosage & administration:****For Tablet :**

**Adult** - 1-2 tablets every 4-6 hours to a maximum of 8 tablets daily.

**For Suspension :****For Children -**

Age less than 3 months : 10 mg/kg body weight, every 3 - 4 times

3 months - 1 year: ½-1 tea-spoonful every 3 - 4 times

1 - 5 years : 1 - 2 tea-spoonful every 3 - 4 times

6 - 12 years : 2 - 4 tea-spoonful every 3 - 4 times

**For Paediatric drops :****For Children -**

0-3 month: 0.5 ml, 4-11 month: 1 ml,

12-13 month: 1.5 ml, 2-3 years: 2 ml,

4-5 years: 3 ml, 4 times daily

**Contraindication:**

Paracetamol is contraindicated in patients with a known hypersensitivity to its component.

**Side effects:**

Side effects are usually hematological reactions that have been reported. Skin eruption have also occurred. Symptoms of overdose includes vomiting, gastrointestinal hemorrhage, liver damage, cerebral edema & renal tubular necrosis. Hypoglycemia has been reported after overdoses of paracetamol.

**Precautions:**

Paracetamol should be given with care to patients with impaired kidney and liver function. It should be also given with care to patients taking other drugs that affects the liver.

**Use in special group:**

**Pregnancy & Lactation:** Though paracetamol appears in breast milk & crosses the placenta, it causes no harmful effects to breast-fed infants or fetus.

**Drug interaction:**

Domperidon & Metoclopramide accelerate the absorption of paracetamol. Colestyramine reduces the absorption of paracetamol. Prolonged regular use of paracetamol possibly enhances anticoagulant effects of coumarine.

**Storage:**

Store in a cool and dry place, protect from light and moisture. Keep all medicines out of the reach of the children.

**Packaging**

**Pamix Tablet:** Each box contains 20 X 10's tablets in blister pack.

**Pamix Suspension:** Each bottle contains 60 ml suspension.

**Pamix Paediatric drops:** Each bottle contains 15 ml drop.

Registered Trade Mark

Manufactured by



**Ziska Pharmaceuticals Ltd.**

Kaliakoir, Gazipur, Bangladesh